Fall Semester 2020
Stat 462 Final Exam
Monday, December 14

Name _____ Student ID #_____

Your exam should consist of 10 problems, followed by a Z-table.

You are allowed a double-sided handwritten note and a calculator.

Problem	Points	S C O R E
1	10	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
10	10	
Total:	100	

1. (10 points) Find the cumulative distribution function for X with probability density function

$$f(x) = \begin{cases} 3x^{-4}, \, x \geq 1; \\ 0, \quad \text{else.} \end{cases}$$

2. (10 points) For $X(t) = 1 + 2t + 1.2W(t)$, find $\mathbb{P}(X(3) > 3 \mid X(1.4) = 0.5)$.

3. (10 points) Find the probability $\mathbb{P}\Big[\max_{0\le t\le 10} W(t) \ge 3,\ -1 \le W(10) \le 4\Big]$.

4. (10 points) Take the continuous-time Markov chain on two states $\{1, 2\}$, with initial distribution $[0.3, 0.7]$ and the generating matrix

$$A = \begin{bmatrix} -2 & 2 \\ 2 & -2 \end{bmatrix}$$

(a) Find the stationary distribution, and the rate of convergence.

(b) Find the distribution at time t.

5. (10 points) Take the compound Poisson process

$$X(t) = \sum_{k=1}^{N(t)} Z_k,$$

where $N = (N(t),\, t \geq 0)$ is the Poisson process with intensity $\lambda = 3$, and $Z_k \sim \text{Exp}(2)$ are identically distributed and independent from N and from each other.

(a) Find the moment generating function, mean, and variance of $X(100)$.

(b) Approximating $X(100)$ by the normal distribution, find the value at risk for confidence level 95%. That is, find the number u such that $\mathbb{P}(X(100) \leq u) = 99\%$.

6. (10 points) For the Poisson process $N = (N(t),\, t \geq 0)$ with intensity $\lambda = 2.5$, find the density, mean, and variance of the time τ_5 when it jumps from 4 to 5.

7. (10 points) Take a discrete-time Markov chain with transition matrix

$$P = \begin{bmatrix} 0.1 & 0.5 & 0.4 \\ 0.2 & 0.8 & 0 \\ 0.7 & 0.2 & 0.1 \end{bmatrix}$$

If the starting point is $X_0 = 2$, find the distribution of X_2.

8. (10 points) For the symmetric random walk with increments ± 1, starting from $S_0 = 0$, find the probability that after 8 steps, it will be at $S_8 = 4$.

9. (10 points) For the symmetric random walk with increments ± 1, starting from $S_0 = 0$, find the probability that after 8 steps, it will be at $S_8 = 4$, but during these 8 steps, it will always stay nonnegative. (Use the reflection principle).

10. (10 points) Find the rate of convergence for a discrete-time Markov chain with transition matrix

$$P = \begin{bmatrix} 0.1 & 0.9 \\ 0.2 & 0.8 \end{bmatrix}$$

Cumulative Probabilities of the Standard Normal Distribution

The table gives the probabilities $\alpha = \Phi(z)$ to the left of given z–values for the standard normal distribution.

For example, the probability that a standard normal random variable Z is less than 1.53 is found at the intersection of the 1.5 rows and the 0.03 column, thus $\Phi(1.53) = P(Z \leq 1.53) = 0.9370$. Due to symmetry it holds $\Phi(-z) = 1 - \Phi(z)$ for all z.

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998

Quantiles of the Standard Normal Distribution

For selected probabilities α, the table shows the values of the quantiles z_α such that $\Phi(z_\alpha) = P(Z \leq z_\alpha) = \alpha$, where Z is a standard normal random variable.

The quantiles satisfy the relation $z_{1-\alpha} = -z_\alpha$.

α	0.9	0.95	0.975	0.99	0.995	0.999
z_α	1.282	1.645	1.960	2.326	2.576	3.090