STAT 461/661
Practice Final 2
December 19, 2018

Name _____ Student ID #_____

TA Name _____ Section Time #_____

Problem	Points	S C O R E
1	10	
2	10	
3	10	
4	10	
5	10	
6	10	
7	10	
8	10	
9	10	
10	10	
Total:	100	

1. (10 points) Take two independent variables $X \sim \text{Exp}(2)$ and $Y \sim \text{Exp}(3)$. Find $\text{Var}\, Z$ for

$$Z = 2X - 3Y + 5.$$

2. (10 points) Take independent random variables $X, Y \sim \text{Uni}[1, 3]$. Find the moment generating function of

$$X - 2Y + 1.$$

3. (10 points) Two random variables X and Y have joint density

$$\begin{cases} \frac{1}{21}(x+y), & 1 \le x \le 3, \quad 0 \le y \le 3; \\ 0, \text{otherwise.} \end{cases}$$

Find the probability $\mathbf{P}(X \ge 2,\ Y \le 2)$.

4. (10 points) An insurance company has $N = 90000$ clients. Each comes with a claim whose distribution has moment generating function

$$F(t) = \frac{1}{3} + \frac{4}{6 - 3t}.$$

The amounts of claims are independent. Find the value-at-risk at confidence level $\alpha = 90\%$.

5. (10 points) There are $N = 1000$ houses, each is on fire independently of other houses with probability $p = 0.4\%$. Find the probability that there are exactly six fires.

6. (10 points) Take a sequence of independent Bernoulli trials, each has probability $p = 0.4$ of success. Let X be the number of trials you need to get to your first success. Find $\mathbf{E}(2X + 3)^2$.

7. (10 points) Find the expectation of a random variable X with CDF

$$F_X(x) = \begin{cases} 0, \ x < -2; \\ 0.4, \ -2 \leq x < -1; \\ 0.7, \ -1 \leq x < 1; \\ 1, \ x \geq 1. \end{cases}$$

8. (10 points) Assume that 0.1% of people from a certain population have a germ. A test gives false positive (that is, shows a person has this germ if this person does not actually have a germ) in 20% of cases when the person does not have this germ. This test gives false negative (shows a person does not have this germ if this person actually has it) in 30% of cases when this person has this germ. Suppose you pick a random person from the population and apply this test twice. Both times it gives you positive result (that is, the test says that this person has this germ). What is the probability that this person actually has this germ?

9. (10 points) Find the distribution of $XY - Z$ for random variables X, Y, Z with joint distribution

X	Y	Z	Prob.
2	1	0	0.5
0	1	-1	0.3
1	0	-2	0.2

10. (10 points) Pick a random number from 1 to 600. Find the probability that this number is divisible by 3 but not by 5.

Cumulative Probabilities of the Standard Normal Distribution

The table gives the probabilities $\alpha = \Phi(z)$ to the left of given z–values for the standard normal distribution.

For example, the probability that a standard normal random variable Z is less than 1.53 is found at the intersection of the 1.5 rows and the 0.03 column, thus $\Phi(1.53) = P(Z \leq 1.53) = 0.9370$. Due to symmetry it holds $\Phi(-z) = 1 - \Phi(z)$ for all z.

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998

Quantiles of the Standard Normal Distribution

For selected probabilities α, the table shows the values of the quantiles z_α such that $\Phi(z_\alpha) = P(Z \leq z_\alpha) = \alpha$, where Z is a standard normal random variable.

The quantiles satisfy the relation $z_{1-\alpha} = -z_\alpha$.

α	0.9	0.95	0.975	0.99	0.995	0.999
z_α	1.282	1.645	1.960	2.326	2.576	3.090